

Schur's Theorem for randomly perturbed sets

Shagnik Das
NTU

Charlotte Knierim
ETH Zürich

Patrick Morris
FU Berlin

Randomly Perturbed (Hyper)graphs
SIAM DM 22 — Pirate Session

17 June 2022

Introduction

An origin story

Theorem (Dirac, 1952)

*If an n -vertex graph G has
minimum degree $\delta(G) \geq \frac{1}{2}n$,
then G is Hamiltonian.*

An origin story

Theorem (Dirac, 1952)

If an n -vertex graph G has minimum degree $\delta(G) \geq \frac{1}{2}n$, then G is Hamiltonian.

Moreover, this bound is best possible.

An origin story

Theorem (Dirac, 1952)

If an n -vertex graph G has minimum degree $\delta(G) \geq \frac{1}{2}n$, then G is Hamiltonian.

Moreover, this bound is best possible.

Theorem (Pósa, 1962)

The random graph $G(n, p)$ is Hamiltonian with probability tending to

0 if it has $o(n \ln n)$ edges.

An origin story

Theorem (Dirac, 1952)

If an n -vertex graph G has minimum degree $\delta(G) \geq \frac{1}{2}n$, then G is Hamiltonian.

Moreover, this bound is best possible.

Theorem (Pósa, 1962)

The random graph $G(n, p)$ is Hamiltonian with probability tending to

- 0 if it has $o(n \ln n)$ edges.*
- 1 if it has $\omega(n \ln n)$ edges.*

An origin story

Theorem (Dirac, 1952)

If an n -vertex graph G has minimum degree $\delta(G) \geq \frac{1}{2}n$, then G is Hamiltonian.

Moreover, this bound is best possible.

Theorem (Pósa, 1962)

The random graph $G(n, p)$ is Hamiltonian with probability tending to

0 if it has $o(n \ln n)$ edges.

1 if it has $\omega(n \ln n)$ edges.

Theorem (Bohman–Frieze–Martin, 2003)

If $\delta(G) = \Omega(n)$, then adding $\Theta(n)$ random edges makes G Hamiltonian with high probability.


Schur's Theorem

Schur triples

A **Schur triple** in $A \subseteq \mathbb{N}$ is a triple $x, y, z \in A$ with $x + y = z$.

Theorem (Schur, 1916)

For every $r \geq 1$, there is an $n = n(r) \in \mathbb{N}$ such that every r -colouring of $[n]$ has a monochromatic Schur triple.




Schur's Theorem

Schur triples

A **Schur triple** in $A \subseteq \mathbb{N}$ is a triple $x, y, z \in A$ with $x + y = z$.

Theorem (Schur, 1916)

For every $r \geq 1$, there is an $n = n(r) \in \mathbb{N}$ such that every r -colouring of $[n]$ has a monochromatic Schur triple.



A Ramsey property

We say a set $A \subseteq \mathbb{N}$ is **r -Schur** if every r -colouring of A has a monochromatic Schur triple.

Schur's Theorem

Schur triples

A **Schur triple** in $A \subseteq \mathbb{N}$ is a triple $x, y, z \in A$ with $x + y = z$.

Theorem (Schur, 1916)

For every $r \geq 1$, there is an $n = n(r) \in \mathbb{N}$ such that every r -colouring of $[n]$ has a monochromatic Schur triple.



A Ramsey property

We say a set $A \subseteq \mathbb{N}$ is **arr-Schur** if every r -colouring of A has a monochromatic Schur triple.

Extremal Schur: sum-free sets

Question

How large can a subset $A \subseteq [n]$ be without being r -Schur?

Extremal Schur: sum-free sets

Question

How large can a subset $A \subseteq [n]$ be without being r -Schur?

$$r = 1$$

Ques: What are the largest sum-free subsets of $[n]$?

Extremal Schur: sum-free sets

Question

How large can a subset $A \subseteq [n]$ be without being r -Schur?

$r = 1$

Ques: What are the largest sum-free subsets of $[n]$?

Ans: Largest sets have size $\frac{n}{2}$

Extremal Schur: sum-free sets

Question

How large can a subset $A \subseteq [n]$ be without being r -Schur?

$r = 1$

Ques: What are the largest sum-free subsets of $[n]$?

Ans: Largest sets have size $\frac{n}{2}$

$$A_{\text{odd}} = \{x \in [n] : x \equiv 1 \pmod{2}\}$$



Extremal Schur: sum-free sets

Question

How large can a subset $A \subseteq [n]$ be without being r -Schur?

$r = 1$

Ques: What are the largest sum-free subsets of $[n]$?

Ans: Largest sets have size $\frac{n}{2}$

$$A_{\text{odd}} = \{x \in [n] : x \equiv 1 \pmod{2}\}$$

$$A_{\text{large}} = \{x \in [n] : x > \frac{n}{2}\}$$



Extremal Schur: more colours

Question

How large can a subset $A \subseteq [n]$ be without being r -Schur?

Extremal Schur: more colours

Question

How large can a subset $A \subseteq [n]$ be without being r -Schur?

Theorem (Hu, 1980)

If $A \subseteq [n]$ with $|A| > \frac{4n}{5}$, then A is 2-Schur.

Extremal Schur: more colours

Question

How large can a subset $A \subseteq [n]$ be without being r -Schur?

Theorem (Hu, 1980)

If $A \subseteq [n]$ with $|A| > \frac{4n}{5}$, then A is 2-Schur.

Bound is tight

$$A = \{x \in [n] : x \equiv 1, 4 \pmod{5}\} \cup \{x \in [n] : x \equiv 2, 3 \pmod{5}\}$$



Extremal Schur: more colours

Question

How large can a subset $A \subseteq [n]$ be without being r -Schur?

Theorem (Hu, 1980)

If $A \subseteq [n]$ with $|A| > \frac{4n}{5}$, then A is 2-Schur.

Bound is tight

$$A = \{x \in [n] : x \equiv 1, 4 \pmod{5}\} \cup \{x \in [n] : x \equiv 2, 3 \pmod{5}\}$$



Even more colours

Problem is open for $r \geq 3$

Extremal Schur: more colours

Question

How large can a subset $A \subseteq [n]$ be without being r -Schur?

Theorem (Hu, 1980)

If $A \subseteq [n]$ with $|A| > \frac{4n}{5}$, then A is 2-Schur.

Bound is tight

$$A = \{x \in [n] : x \equiv 1, 4 \pmod{5}\} \cup \{x \in [n] : x \equiv 2, 3 \pmod{5}\}$$



Even more colours

Problem is open for $r \geq 3$

- Conjecture from Abbott and Wang (1977)

Probabilistic Schur: sum-free sets

Definition (Binomial random integer subsets)

Given $p \in [0, 1]$, choose each element of $[n]$ to be in $[n]_p$ independently with probability p .

Probabilistic Schur: sum-free sets

Definition (Binomial random integer subsets)

Given $p \in [0, 1]$, choose each element of $[n]$ to be in $[n]_p$ independently with probability p .

Question

At what density p is $[n]_p \subseteq [n]$ typically r -Schur?

Probabilistic Schur: sum-free sets

Definition (Binomial random integer subsets)

Given $p \in [0, 1]$, choose each element of $[n]$ to be in $[n]_p$ independently with probability p .

Question

At what density p is $[n]_p \subseteq [n]$ typically r -Schur?

Proposition ($r = 1$)

- (0) If $p = o(n^{-2/3})$, then w.h.p. $[n]_p$ is sum-free.
- (1) If $p = \omega(n^{-2/3})$, then w.h.p. $[n]_p$ is not sum-free.

Probabilistic Schur: more colours

Proposition ($r = 1$)

- (0) If $p = o(n^{-2/3})$, then w.h.p. $[n]_p$ is sum-free.
- (1) If $p = \omega(n^{-2/3})$, then w.h.p. $[n]_p$ is not sum-free.

Probabilistic Schur: more colours

Proposition ($r = 1$)

- (0) If $p = o(n^{-2/3})$, then w.h.p. $[n]_p$ is sum-free.
- (1) If $p = \omega(n^{-2/3})$, then w.h.p. $[n]_p$ is not sum-free.

Theorem (Graham–Rödl–Ruciński, 1996)

For $r = 2$, the following hold.

- (0) If $p = o(n^{-1/2})$, then w.h.p. $[n]_p$ is not r -Schur.
- (1) If $p = \omega(n^{-1/2})$, then w.h.p. $[n]_p$ is r -Schur.

Probabilistic Schur: more colours

Proposition ($r = 1$)

- (0) If $p = o(n^{-2/3})$, then w.h.p. $[n]_p$ is sum-free.
- (1) If $p = \omega(n^{-2/3})$, then w.h.p. $[n]_p$ is not sum-free.

Theorem (Graham–Rödl–Ruciński, 1996)

For $r = 2$, the following hold.

- (0) If $p = o(n^{-1/2})$, then w.h.p. $[n]_p$ is not r -Schur.
- (1) If $p = \omega(n^{-1/2})$, then w.h.p. $[n]_p$ is r -Schur.

Theorem (Rödl–Ruciński, 1997)

For $r \geq 3$, the following hold.

- (0) If $p = o(n^{-1/2})$, then w.h.p. $[n]_p$ is not r -Schur.
- (1) If $p = \omega(n^{-1/2})$, then w.h.p. $[n]_p$ is r -Schur.

Probabilistic Schur: more colours

Proposition ($r = 1$)

- (0) If $p = o(n^{-2/3})$, then w.h.p. $[n]_p$ is sum-free.
- (1) If $p = \omega(n^{-2/3})$, then w.h.p. $[n]_p$ is not sum-free.

Theorem (Graham–Rödl–Ruciński, 1996)

For $r = 2$, the following hold.

- (0) If $p = o(n^{-1/2})$, then w.h.p. $[n]_p$ is not r -Schur.
- (1) If $p = \omega(n^{-1/2})$, then w.h.p. $[n]_p$ is r -Schur.

Theorem (Rödl–Ruciński, 1997)

For $r \geq 3$, the following hold.

- (0) If $p = o(n^{-1/2})$, then w.h.p. $[n]_p$ is not r -Schur.
- (1) If $p = \omega(n^{-1/2})$, then w.h.p. $[n]_p$ is r -Schur.

Perturbed Schur

Shaken, not Schurred

Given $A \subseteq [n]$, for which p will $A \cup [n]_p$ be 2-Schur w.h.p.?



Perturbed Schur

Shaken, not Schurred

Given $A \subseteq [n]$, for which p will $A \cup [n]_p$ be 2-Schur w.h.p.?



Observation

- (*Extremal Schur*) If $|A| > \frac{4}{5}n$, we can take $p = 0$

Perturbed Schur

Shaken, not Schurred

Given $A \subseteq [n]$, for which p will $A \cup [n]_p$ be 2-Schur w.h.p.?



Observation

- (*Extremal Schur*) If $|A| > \frac{4}{5}n$, we can take $p = 0$
- (*Probabilistic Schur*) If $|A| = 0$, we need $p = \Omega(n^{-1/2})$

Perturbed Schur

Shaken, not Schurred

Given $A \subseteq [n]$, for which p will $A \cup [n]_p$ be 2-Schur w.h.p.?



Observation

- (Extremal Schur) If $|A| > \frac{4}{5}n$, we can take $p = 0$
- (Probabilistic Schur) If $|A| = 0$, we need $p = \Omega(n^{-1/2})$

Theorem (Aigner-Horev–Person, 2019)

If $|A| = \Omega(n)$ and $p = \omega(n^{-2/3})$, then $A \cup [n]_p$ is 2-Schur w.h.p.

Perturbed Schur

Shaken, not Schurred

Given $A \subseteq [n]$, for which p will $A \cup [n]_p$ be 2-Schur w.h.p.?



Observation

- (Extremal Schur) If $|A| > \frac{4}{5}n$, we can take $p = 0$
- (Probabilistic Schur) If $|A| = 0$, we need $p = \Omega(n^{-1/2})$


Theorem (Aigner-Horev–Person, 2019)

If $|A| = \Omega(n)$ and $p = \omega(n^{-2/3})$, then $A \cup [n]_p$ is 2-Schur w.h.p.

Observation

Result is best possible for $|A| \leq \frac{n}{2}$: take A sum-free, and $A \cup [n]_p$

A pictorial summary



Our results

Denser sets

We complete the picture for sets $|A| > \frac{n}{2}$

Denser sets

We complete the picture for sets $|A| > \frac{n}{2}$

Theorem (D.–Knierim–Morris, 2022+)

For $n, t \in \mathbb{N}$ with $\frac{n}{2} + t \leq \frac{4n}{5}$, define $p(n, t) = \min \{n^{-2/3}, t^{-1}\}$.

Denser sets

We complete the picture for sets $|A| > \frac{n}{2}$

Theorem (D.–Knierim–Morris, 2022+)

For $n, t \in \mathbb{N}$ with $\frac{n}{2} + t \leq \frac{4n}{5}$, define $p(n, t) = \min \{n^{-2/3}, t^{-1}\}$.

(0) $\exists A \subset [n]$, $|A| = \frac{n}{2} + t$, such that if $p = o(p(n, t))$, then $A \cup [n]_p$ is w.h.p. not 2-Schur.

Denser sets

We complete the picture for sets $|A| > \frac{n}{2}$

Theorem (D.–Knierim–Morris, 2022+)

For $n, t \in \mathbb{N}$ with $\frac{n}{2} + t \leq \frac{4n}{5}$, define $p(n, t) = \min \{n^{-2/3}, t^{-1}\}$.

- (0) $\exists A \subset [n]$, $|A| = \frac{n}{2} + t$, such that if $p = o(p(n, t))$, then $A \cup [n]_p$ is w.h.p. not 2-Schur.
- (1) $\forall A \subset [n]$, $|A| = \frac{n}{2} + t$, and $p = \omega(p(n, t))$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Denser sets

We complete the picture for sets $|A| > \frac{n}{2}$

Theorem (D.–Knierim–Morris, 2022+)

For $n, t \in \mathbb{N}$ with $\frac{n}{2} + t \leq \frac{4n}{5}$, define $p(n, t) = \min \{n^{-2/3}, t^{-1}\}$.

(0) $\exists A \subset [n]$, $|A| = \frac{n}{2} + t$, such that if $p = o(p(n, t))$, then $A \cup [n]_p$ is w.h.p. not 2-Schur.

(1) $\forall A \subset [n]$, $|A| = \frac{n}{2} + t$, and $p = \omega(p(n, t))$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Observation

We need $t = \omega(n^{2/3})$ before we save any further randomness.

Perturbed stability

Theorem (D.–Knierim–Morris, 2022+)

If $A \subset [n]$ with $|A| = \frac{n}{2} + t$, and $q = \omega(n^{-1})$ is such that $A \cup [n]_q$ is w.h.p. not 2-Schur, then either $|A_{\text{large}} \setminus A| = O(q^{-1})$ or $|A_{\text{odd}} \setminus A| = O(q^{-2}n^{-1})$.

Perturbed stability

Theorem (D.–Knierim–Morris, 2022+)

If $A \subset [n]$ with $|A| = \frac{n}{2} + t$, and $q = \omega(n^{-1})$ is such that $A \cup [n]_q$ is w.h.p. not 2-Schur, then either $|A_{\text{large}} \setminus A| = O(q^{-1})$ or $|A_{\text{odd}} \setminus A| = O(q^{-2}n^{-1})$.

Remark

- *If A is not close to an extremal sum-free set, then very little randomness is needed*

Perturbed stability

Theorem (D.–Knierim–Morris, 2022+)

If $A \subset [n]$ with $|A| = \frac{n}{2} + t$, and $q = \omega(n^{-1})$ is such that $A \cup [n]_q$ is w.h.p. not 2-Schur, then either $|A_{\text{large}} \setminus A| = O(q^{-1})$ or $|A_{\text{odd}} \setminus A| = O(q^{-2}n^{-1})$.

Remark

- If A is not close to an extremal sum-free set, then very little randomness is needed
- A must contain at least t even numbers
 \Rightarrow first case cannot occur if $q = \omega((nt)^{-1/2})$

Perturbed stability

Theorem (D.–Knierim–Morris, 2022+)

If $A \subset [n]$ with $|A| = \frac{n}{2} + t$, and $q = \omega(n^{-1})$ is such that $A \cup [n]_q$ is w.h.p. not 2-Schur, then either $|A_{\text{large}} \setminus A| = O(q^{-1})$ or $|A_{\text{odd}} \setminus A| = O(q^{-2}n^{-1})$.

Remark

- *If A is not close to an extremal sum-free set, then very little randomness is needed*
- *A must contain at least t even numbers*
 - \Rightarrow *first case cannot occur if $q = \omega((nt)^{-1/2})$*
 - *when $n^{1/3} \ll t \ll n$, we have $(nt)^{-1/2} \ll \min\{n^{-2/3}, t^{-1}\}$*

Perturbed stability


Theorem (D.–Knierim–Morris, 2022+)

If $A \subset [n]$ with $|A| = \frac{n}{2} + t$, and $q = \omega(n^{-1})$ is such that $A \cup [n]_q$ is w.h.p. not 2-Schur, then either $|A_{\text{large}} \setminus A| = O(q^{-1})$ or $|A_{\text{odd}} \setminus A| = O(q^{-2}n^{-1})$.

Remark

- If A is not close to an extremal sum-free set, then very little randomness is needed
- A must contain at least t even numbers
 - \Rightarrow first case cannot occur if $q = \omega((nt)^{-1/2})$
 - when $n^{1/3} \ll t \ll n$, we have $(nt)^{-1/2} \ll \min\{n^{-2/3}, t^{-1}\}$
 - $\Rightarrow A_{\text{large}}$ “more sum-free” than A_{odd}

An updated picture



The sparse range

Question

What happens when our initial set is sparse, $|A| = o(n)$?

The sparse range

Question

What happens when our initial set is sparse, $|A| = o(n)$?

Observation

No gain if $|A| = o(n^{1/2})$:

The sparse range

Question

What happens when our initial set is sparse, $|A| = o(n)$?

Observation

No gain if $|A| = o(n^{1/2})$:

Let $d = \frac{|A|}{n}$ and take $A \sim [n]_d$

The sparse range

Question

What happens when our initial set is sparse, $|A| = o(n)$?

Observation

No gain if $|A| = o(n^{1/2})$:

Let $d = \frac{|A|}{n}$ and take $A \sim [n]_d$

$\Rightarrow A \cup [n]_p \sim [n]_q$ for $q \approx d + p$

The sparse range

Question

What happens when our initial set is sparse, $|A| = o(n)$?

Observation

No gain if $|A| = o(n^{1/2})$:

Let $d = \frac{|A|}{n}$ and take $A \sim [n]_d$

$\Rightarrow A \cup [n]_p \sim [n]_q$ for $q \approx d + p$

\Rightarrow not 2-Schur unless $p = \Omega(n^{-1/2})$

The sparse range

Question

What happens when our initial set is sparse, $|A| = o(n)$?

Observation

No gain if $|A| = o(n^{1/2})$:

Let $d = \frac{|A|}{n}$ and take $A \sim [n]_d$

$\Rightarrow A \cup [n]_p \sim [n]_q$ for $q \approx d + p$

\Rightarrow not 2-Schur unless $p = \Omega(n^{-1/2})$

Question

*What happens when our initial set is moderately sparse,
 $\Omega(n^{1/2}) = |A| = o(n)$?*

Sparse results

Theorem (D.–Knierim–Morris, 2022+)

Let $n, s \in \mathbb{N}$ satisfy $\Omega(n^{1/2}) = s \leq \frac{n}{2}$.

Sparse results

Theorem (D.–Knierim–Morris, 2022+)

Let $n, s \in \mathbb{N}$ satisfy $\Omega(n^{1/2}) = s \leq \frac{n}{2}$.

(0) $\exists A \subset [n]$, $|A| = s$, such that if $p = o((ns)^{-1/3})$, then $A \cup [n]_p$ is w.h.p. not 2-Schur.

Sparse results

Theorem (D.–Knierim–Morris, 2022+)

Let $n, s \in \mathbb{N}$ satisfy $\Omega(n^{1/2}) = s \leq \frac{n}{2}$.

- (0) $\exists A \subset [n]$, $|A| = s$, such that if $p = o((ns)^{-1/3})$, then $A \cup [n]_p$ is w.h.p. not 2-Schur.
- (1) $\forall A \subset [n]$, $|A| = s$, and $p = \tilde{\omega}((n^{13}s)^{-1/27})$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Sparse results

Theorem (D.–Knierim–Morris, 2022+)

Let $n, s \in \mathbb{N}$ satisfy $\Omega(n^{1/2}) = s \leq \frac{n}{2}$.

- (0) $\exists A \subset [n]$, $|A| = s$, such that if $p = o((ns)^{-1/3})$, then $A \cup [n]_p$ is w.h.p. not 2-Schur.
- (1) $\forall A \subset [n]$, $|A| = s$, and $p = \tilde{\omega}((n^{13}s)^{-1/27})$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Remark

- Lower bound interpolates between $n^{-1/2}$ and $n^{-2/3}$

Sparse results

Theorem (D.–Knierim–Morris, 2022+)

Let $n, s \in \mathbb{N}$ satisfy $\Omega(n^{1/2}) = s \leq \frac{n}{2}$.

- (0) $\exists A \subset [n]$, $|A| = s$, such that if $p = o((ns)^{-1/3})$, then $A \cup [n]_p$ is w.h.p. not 2-Schur.
- (1) $\forall A \subset [n]$, $|A| = s$, and $p = \tilde{\omega}((n^{13}s)^{-1/27})$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Remark

- Lower bound interpolates between $n^{-1/2}$ and $n^{-2/3}$
- Upper bound is $o(n^{-1/2})$ when $s = \tilde{\omega}(n^{1/2})$

Proof sketches

Lower bound

Theorem (D.–Knierim–Morris, 2022+)

(0) $\exists A \subset [n]$, $|A| = s$, such that if $p = o((ns)^{-1/3})$, then $A \cup [n]_p$ is w.h.p. not 2-Schur.

Lower bound

Theorem (D.–Knierim–Morris, 2022+)

(0) $\exists A \subset [n]$, $|A| = s$, such that if $p = o((ns)^{-1/3})$, then $A \cup [n]_p$ is w.h.p. not 2-Schur.

Proof idea

- Take $A = [n - s + 1, n]$, coloured green

Lower bound

Theorem (D.–Knierim–Morris, 2022+)

(0) $\exists A \subset [n]$, $|A| = s$, such that if $p = o((ns)^{-1/3})$, then $A \cup [n]_p$ is w.h.p. not 2-Schur.

Proof idea

- Take $A = [n - s + 1, n]$, coloured green
- Algorithmic recolouring of random elements
(à la random Ramsey lower bounds)

Lower bound

Theorem (D.–Knierim–Morris, 2022+)

(0) $\exists A \subset [n]$, $|A| = s$, such that if $p = o((ns)^{-1/3})$, then $A \cup [n]_p$ is w.h.p. not 2-Schur.

Proof idea

- Take $A = [n - s + 1, n]$, coloured green
- Algorithmic recolouring of random elements
(à la random Ramsey lower bounds)
 - ▶ if no good colouring exists \Rightarrow cycle of Schur triples

Lower bound

Theorem (D.–Knierim–Morris, 2022+)

(0) $\exists A \subset [n]$, $|A| = s$, such that if $p = o((ns)^{-1/3})$, then $A \cup [n]_p$ is w.h.p. not 2-Schur.

Proof idea

- Take $A = [n - s + 1, n]$, coloured green
- Algorithmic recolouring of random elements
(à la random Ramsey lower bounds)
 - ▶ if no good colouring exists \Rightarrow cycle of Schur triples
 - ▶ Such a cycle unlikely to exist in random elements



Upper bound

Theorem (D.–Knierim–Morris, 2022+)

- (1) $\forall A \subset [n]$, $|A| = s$, and $p = \tilde{\omega}((n^{13}s)^{-1/27})$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Upper bound

Theorem (D.–Knierim–Morris, 2022+)

(1) $\forall A \subset [n]$, $|A| = s$, and $p = \tilde{\omega}((n^{13}s)^{-1/27})$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Proof idea

- Show the random set $[n]_p$ is incompatible with any colouring

Upper bound

Theorem (D.–Knierim–Morris, 2022+)

(1) $\forall A \subset [n]$, $|A| = s$, and $p = \tilde{\omega}((n^{13}s)^{-1/27})$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Proof idea

- Show the random set $[n]_p$ is incompatible with any colouring
- Problem: too many colourings for a union bound

Upper bound

Theorem (D.–Knierim–Morris, 2022+)

(1) $\forall A \subset [n]$, $|A| = s$, and $p = \tilde{\omega}((n^{13}s)^{-1/27})$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Proof idea

- Show the random set $[n]_p$ is incompatible with any colouring
- Problem: too many colourings for a union bound
- Solution: build a hypergraph

Upper bound

Theorem (D.–Knierim–Morris, 2022+)

(1) $\forall A \subset [n]$, $|A| = s$, and $p = \tilde{\omega}((n^{13}s)^{-1/27})$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Proof idea

- Show the random set $[n]_p$ is incompatible with any colouring
- Problem: too many colourings for a union bound
- Solution: build a hypergraph
 - ▶ vertices \leftrightarrow coloured elements

Upper bound

Theorem (D.–Knierim–Morris, 2022+)

(1) $\forall A \subset [n]$, $|A| = s$, and $p = \tilde{\omega}((n^{13}s)^{-1/27})$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Proof idea

- Show the random set $[n]_p$ is incompatible with any colouring
- Problem: too many colourings for a union bound
- Solution: build a hypergraph
 - ▶ vertices \leftrightarrow coloured elements
 - ▶ edges \leftrightarrow small configurations forcing mc Schur triples

Upper bound

Theorem (D.–Knierim–Morris, 2022+)

(1) $\forall A \subset [n]$, $|A| = s$, and $p = \tilde{\omega}((n^{13}s)^{-1/27})$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Proof idea

- Show the random set $[n]_p$ is incompatible with any colouring
 - Problem: too many colourings for a union bound
 - Solution: build a hypergraph
 - ▶ vertices \leftrightarrow coloured elements
 - ▶ edges \leftrightarrow small configurations forcing mc Schur triples
- \Rightarrow mc-Schur-triple-free colourings \rightarrow independent sets

Upper bound

Theorem (D.–Knierim–Morris, 2022+)

(1) $\forall A \subset [n]$, $|A| = s$, and $p = \tilde{\omega}((n^{13}s)^{-1/27})$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Proof idea

- Show the random set $[n]_p$ is incompatible with any colouring
- Problem: too many colourings for a union bound
- Solution: build a hypergraph
 - ▶ vertices \leftrightarrow coloured elements
 - ▶ edges \leftrightarrow small configurations forcing mc Schur triples
 - \Rightarrow mc-Schur-triple-free colourings \rightarrow independent sets
- Hypergraph containers: can group together similar colourings

Upper bound

Theorem (D.–Knierim–Morris, 2022+)

(1) $\forall A \subset [n]$, $|A| = s$, and $p = \tilde{\omega}((n^{13}s)^{-1/27})$, $A \cup [n]_p$ is w.h.p. 2-Schur.

Proof idea

- Show the random set $[n]_p$ is incompatible with any colouring
- Problem: too many colourings for a union bound
- Solution: build a hypergraph
 - ▶ vertices \leftrightarrow coloured elements
 - ▶ edges \leftrightarrow small configurations forcing mc Schur triples
 - \Rightarrow mc-Schur-triple-free colourings \rightarrow independent sets
- Hypergraph containers: can group together similar colourings
 - ▶ apply union bound more efficiently



The colouring hypergraph — vertices

Definition (Colouring hypergraph)

Given $A \subseteq [n]$, $|A| = s$, the colouring hypergraph \mathcal{H}_A has vertices $V(\mathcal{H}_A) = V_R \cup V_G$, which are two disjoint copies of $[n]$.

The colouring hypergraph — vertices

Definition (Colouring hypergraph)

Given $A \subseteq [n]$, $|A| = s$, the colouring hypergraph \mathcal{H}_A has vertices $V(\mathcal{H}_A) = V_R \cup V_G$, which are two disjoint copies of $[n]$.

Modelling colourings

Given any $S \subseteq [n]$, map colourings $\varphi : S \rightarrow \{\text{red}, \text{green}\}$ to

$$\{i \in V_R : i \in S, \varphi(i) = \text{red}\} \cup \{i \in V_B : i \in S, \varphi(i) = \text{green}\} \subseteq V(\mathcal{H}_A).$$

The colouring hypergraph — vertices

Definition (Colouring hypergraph)

Given $A \subseteq [n]$, $|A| = s$, the colouring hypergraph \mathcal{H}_A has vertices $V(\mathcal{H}_A) = V_R \cup V_G$, which are two disjoint copies of $[n]$.

Modelling colourings

Given any $S \subseteq [n]$, map colourings $\varphi : S \rightarrow \{\text{red}, \text{green}\}$ to

$$\{i \in V_R : i \in S, \varphi(i) = \text{red}\} \cup \{i \in V_B : i \in S, \varphi(i) = \text{green}\} \subseteq V(\mathcal{H}_A).$$

V_R 1 2 3 4 5 6 7 8 9 10

V_G 1 2 3 4 5 6 7 8 9 10

The colouring hypergraph — vertices

Definition (Colouring hypergraph)

Given $A \subseteq [n]$, $|A| = s$, the colouring hypergraph \mathcal{H}_A has vertices $V(\mathcal{H}_A) = V_R \cup V_G$, which are two disjoint copies of $[n]$.

Modelling colourings

Given any $S \subseteq [n]$, map colourings $\varphi : S \rightarrow \{\text{red}, \text{green}\}$ to

$$\{i \in V_R : i \in S, \varphi(i) = \text{red}\} \cup \{i \in V_B : i \in S, \varphi(i) = \text{green}\} \subseteq V(\mathcal{H}_A).$$

V_R	1	2	3	4	5	6	7	8	9	10
V_G	1	2	3	4	5	6	7	8	9	10
$\varphi(S)$	2	5	6	8	10					

The colouring hypergraph — edges

Definition (Hypergraph edges)

For every $a \in A$ and $x, y, z, w \in [n]$ such that a, x, y and a, z, w form Schur triples, we add a 4-edge on the vertices $x, y \in V_R$ and $z, w \in V_G$.

The colouring hypergraph — edges

Definition (Hypergraph edges)

For every $a \in A$ and $x, y, z, w \in [n]$ such that a, x, y and a, z, w form Schur triples, we add a 4-edge on the vertices $x, y \in V_R$ and $z, w \in V_G$.

Observation

- If x, y are coloured red and z, w are coloured green, we cannot colour $a \in A$

The colouring hypergraph — edges

Definition (Hypergraph edges)

For every $a \in A$ and $x, y, z, w \in [n]$ such that a, x, y and a, z, w form Schur triples, we add a 4-edge on the vertices $x, y \in V_R$ and $z, w \in V_G$.

Observation

- If x, y are coloured red and z, w are coloured green, we cannot colour $a \in A$

\Rightarrow Any Schur colouring of $A \cup [n]_p$ is independent in \mathcal{H}_A

Containers for colourings

Proposition

For every $\varepsilon > 0$ there is some $c = c_\varepsilon$ such that, if $A \subseteq [n]$ is of size $s = \Omega(n^{1/2})$, then there is a collection \mathcal{C} of subsets of $V(\mathcal{H}_A)$ for which:

1. For every $P \subseteq [n]$ and Schur colouring φ of $A \cup P$, there is some $C \in \mathcal{C}$ such that $\varphi \subseteq C$.
2. For every $C \in \mathcal{C}$, $e(\mathcal{H}_A[C]) \leq \varepsilon s n^2$.
3. $\log |\mathcal{C}| \leq c s^{-1/3} n^{2/3} \log n$.

Containers for colourings

Proposition

For every $\varepsilon > 0$ there is some $c = c_\varepsilon$ such that, if $A \subseteq [n]$ is of size $s = \Omega(n^{1/2})$, then there is a collection \mathcal{C} of subsets of $V(\mathcal{H}_A)$ for which:

1. For every $P \subseteq [n]$ and Schur colouring φ of $A \cup P$, there is some $C \in \mathcal{C}$ such that $\varphi \subseteq C$.
2. For every $C \in \mathcal{C}$, $e(\mathcal{H}_A[C]) \leq \varepsilon s n^2$.
3. $\log |\mathcal{C}| \leq c s^{-1/3} n^{2/3} \log n$.

Goal

Show that for every container $C \in \mathcal{C}$, the probability that $P \sim [n]_p$ admits a Schur colouring φ of $A \cup P$ with $\varphi \subseteq C$ is very small.

Anatomy of a container

Partition

Given $C \subseteq V(\mathcal{H}_A)$, we can partition $[n]$ into four sets:

Anatomy of a container

Partition

Given $C \subseteq V(\mathcal{H}_A)$, we can partition $[n]$ into four sets:

- missing elements $M_C: i \notin C \cap V_R, C \cap V_G$

Anatomy of a container

Partition

Given $C \subseteq V(\mathcal{H}_A)$, we can partition $[n]$ into four sets:

- missing elements $M_C: i \notin C \cap V_R, C \cap V_G$
- red elements $R_C: i \in C \cap V_R, i \notin C \cap V_G$

Anatomy of a container

Partition

Given $C \subseteq V(\mathcal{H}_A)$, we can partition $[n]$ into four sets:

- missing elements $M_C: i \notin C \cap V_R, C \cap V_G$
- red elements $R_C: i \in C \cap V_R, i \notin C \cap V_G$
- green elements $G_C: i \notin C \cap V_R, i \in C \cap V_G$

Anatomy of a container

Partition

Given $C \subseteq V(\mathcal{H}_A)$, we can partition $[n]$ into four sets:

- missing elements $M_C: i \notin C \cap V_R, C \cap V_G$
- red elements $R_C: i \in C \cap V_R, i \notin C \cap V_G$
- green elements $G_C: i \notin C \cap V_R, i \in C \cap V_G$
- two-coloured elements $T_C: i \in C \cap V_R, C \cap V_G$

Anatomy of a container

Partition

Given $C \subseteq V(\mathcal{H}_A)$, we can partition $[n]$ into four sets:

- missing elements M_C : $i \notin C \cap V_R, C \cap V_G$
- red elements R_C : $i \in C \cap V_R, i \notin C \cap V_G$
- green elements G_C : $i \notin C \cap V_R, i \in C \cap V_G$
- two-coloured elements T_C : $i \in C \cap V_R, C \cap V_G$

Observation

- C prescribes the colours of the elements in R_C and G_C
- Elements from M_C cannot be coloured

Classifying containers

There are three types of containers in the world

Classifying containers

There are three types of containers in the world

Type I

Linearly many elements are missing: $|M_C| \geq \varepsilon n$

\Rightarrow very unlikely that all of them will be missing in $P \sim [n]_p$

Classifying containers

There are three types of containers in the world

Type I

Linearly many elements are missing: $|M_C| \geq \varepsilon n$

\Rightarrow very unlikely that all of them will be missing in $P \sim [n]_p$

Type II

Quadratically many Schur triples in R_C or G_C

\Rightarrow (Janson) whp such a triple appears in $P \sim [n]_p \rightarrow \text{mc}$

Classifying containers

There are three types of containers in the world

Type I

Linearly many elements are missing: $|M_C| \geq \varepsilon n$

\Rightarrow very unlikely that all of them will be missing in $P \sim [n]_p$

Type II

Quadratically many Schur triples in R_C or G_C

\Rightarrow (Janson) whp such a triple appears in $P \sim [n]_p \rightarrow \text{mc}$

Type III

None of the above

Containers of the third type

Ingredients

- Almost all elements in $[n]$ receive at least one colour
- Very few Schur triples in R_C, G_C
- $\mathcal{H}_A[C]$ has very few edges

Containers of the third type

Ingredients

- Almost all elements in $[n]$ receive at least one colour
- Very few Schur triples in R_C, G_C
- $\mathcal{H}_A[C]$ has very few edges

Recipe

Green's Removal Lemma + Stability + Janson

Containers of the third type

Ingredients

- Almost all elements in $[n]$ receive at least one colour
- Very few Schur triples in R_C, G_C
- $\mathcal{H}_A[C]$ has very few edges

Recipe

Green's Removal Lemma + Stability + Janson

Outcome

$P \sim [n]_p$ almost surely
contains a wicket

Containers of the third type

Ingredients

- Almost all elements in $[n]$ receive at least one colour
- Very few Schur triples in R_C, G_C
- $\mathcal{H}_A[C]$ has very few edges

Recipe

Green's Removal Lemma + Stability + Janson

Outcome

$P \sim [n]_p$ almost surely
contains a wicket

$$\begin{array}{ccccc} x_1 & + & x_2 & = & x_3 \\ + & & + & & + \\ y_1 & & y_2 & & y_3 \\ || & & || & & || \\ z_1 & & z_2 & & z_3 \end{array}$$

Containers of the third type

Ingredients

- Almost all elements in $[n]$ receive at least one colour
- Very few Schur triples in R_C, G_C
- $\mathcal{H}_A[C]$ has very few edges

Recipe

Green's Removal Lemma + Stability + Janson

Outcome


$P \sim [n]_p$ almost surely
contains a wicket

No way to colour x_1, x_2, x_3


x_1	+	x_2	=	x_3
+		+		+
y_1		y_2		y_3
z_1		z_2		z_3

Concluding remarks

Progress made



Progress made



Open problems

The sparse range

Is $p = (ns)^{-1/3}$ the correct threshold for sets of size s ?

Open problems

The sparse range

Is $p = (ns)^{-1/3}$ the correct threshold for sets of size s ?

More colours

What is the threshold for making sets r -Schur, for $r \geq 3$?

Open problems

The sparse range

Is $p = (ns)^{-1/3}$ the correct threshold for sets of size s ?

More colours

What is the threshold for making sets r -Schur, for $r \geq 3$?

Other systems

Randomly perturbed version of Van der Waerden's theorem?

Open problems

The sparse range

Is $p = (ns)^{-1/3}$ the correct threshold for sets of size s ?

More colours

What is the threshold for making sets r -Schur, for $r \geq 3$?

Other systems

Randomly perturbed version of Van der Waerden's theorem?

Thank you for
your attention!